

# Math 123: Volumes and Arc Length

Ryan Blair

CSU Long Beach

Tuesday February 23, 2016

# Outline

1 Volumes of Rotation

2 Arc Length

# Volumes of solids of rotation

Replace all  $x$ 's with  $y$ 's in the following formulas to get other valid expressions for volume.

**Disks:**

$$\text{Vol} = \int_a^b \pi(\text{radius in terms of } x)^2 dx$$

**Shells:**

$$\text{Vol} = \int_a^b 2\pi(\text{radius in terms of } x)(\text{height in terms of } x) dx$$

**Washers:**

$$\text{Vol} = \int_a^b \pi(\text{outer radius in terms of } x)^2 - \pi(\text{inner radius in terms of } x)^2 dx$$

# Volumes of solids of rotation

Replace all  $x$ 's with  $y$ 's in the following formulas to get other valid expressions for volume.

**Disks:**

$$\text{Vol} = \int_a^b \pi(\text{radius in terms of } x)^2 dx$$

**Shells:**

$$\text{Vol} = \int_a^b 2\pi(\text{radius in terms of } x)(\text{height in terms of } x) dx$$

**Washers:**

$$\text{Vol} = \int_a^b \pi(\text{outer radius in terms of } x)^2 - \pi(\text{inner radius in terms of } x)^2 dx$$

**Exercise:** Find the volume of the object obtained by rotating the region bounded by  $y = 2x^2 - x^3$  and  $y = 0$  about the  $y$ -axis.

# Volumes of solids of rotation

Replace all  $x$ 's with  $y$ 's in the following formulas to get other valid expressions for volume.

## Disks:

$$\text{Vol} = \int_a^b \pi(\text{radius in terms of } x)^2 dx$$

## Shells:

$$\text{Vol} = \int_a^b 2\pi(\text{radius in terms of } x)(\text{height in terms of } x) dx$$

## Washers:

$$\text{Vol} = \int_a^b \pi(\text{outer radius in terms of } x)^2 - \pi(\text{inner radius in terms of } x)^2 dx$$

**Exercise:** Find the volume of the object obtained by rotating the region bounded by the curves  $y = \cos(x) + 1$ ,  $y = 0$  and  $x = 0$  that contains  $(1, 1)$  about the  $x$ -axis.

# Volumes of solids of rotation

Replace all  $x$ 's with  $y$ 's in the following formulas to get other valid expressions for volume.

## Disks:

$$\text{Vol} = \int_a^b \pi(\text{radius in terms of } x)^2 dx$$

## Shells:

$$\text{Vol} = \int_a^b 2\pi(\text{radius in terms of } x)(\text{height in terms of } x) dx$$

## Washers:

$$\text{Vol} = \int_a^b \pi(\text{outer radius in terms of } x)^2 - \pi(\text{inner radius in terms of } x)^2 dx$$

**Exercise:** Find the volume of the object obtained by rotating the region bounded by the curves  $y = \cos(x) + 1$ ,  $y = 0$  and  $x = 0$  that contains  $(1, 1)$  about the  $y$ -axis.

# The length of a curve

Lets find the length of a curve by approximating by line segments.

# The length of a curve

Lets find the length of a curve by approximating by line segments.

If  $f$  is continuous on the interval  $[a, b]$ , then the length of the graph of  $f$  from  $a$  to  $b$  is

$$L = \int_a^b \sqrt{1 + (f'(x))^2}$$



# The length of a curve

Lets find the length of a curve by approximating by line segments.

If  $f$  is continuous on the interval  $[a, b]$ , then the length of the graph of  $f$  from  $a$  to  $b$  is

$$L = \int_a^b \sqrt{1 + (f'(x))^2}$$

Example: Find circumference of the circle  $x^2 + y^2 = 4$ .

# The length of a curve

Lets find the length of a curve by approximating by line segments.

If  $f$  is continuous on the interval  $[a, b]$ , then the length of the graph of  $f$  from  $a$  to  $b$  is

$$L = \int_a^b \sqrt{1 + (f'(x))^2}$$

Example: Find circumference of the circle  $x^2 + y^2 = 4$ .

Example: Find the length of the curve  $y = \ln(\cos(x))$  between  $x = 0$  and  $x = \frac{\pi}{3}$