


## Problems

LO1. Answer the following.

- Provide the definition of what it means to be a mapping reduction from problem  $A$  to problem  $B$ . Hint: do *not* assume that  $A$  and  $B$  are decision problems.
- The simple graph  $G = (V, E)$  shown below is an instance of the **Maximum Independent Set (MIS)** optimization problem. Draw  $f(G)$ , where  $f$  is the mapping reduction from MIS to **Maximum Clique** provided in lecture.



- Verify that  $f$  is valid for input  $G$  in the sense that both  $G$  and  $f(G)$  have the same solution. Hint: for both problems please assume that a “solution” is represented by a subset of vertices.

LO2. An instance of the **Dominating Set** decision problem is a pair  $(G, k)$ , where  $G = (V, E)$  is a simple graph,  $k \geq 0$  is an integer, and the problem is to decide if  $G$  has a **dominating set** of size  $k$ , i.e. a set  $D \subseteq V$  for which every vertex  $v \in V$  is adjacent to at least one vertex in  $D$ . For example, for the graph  $G$  shown above,  $(G, k = 2)$  is a positive instance of **Dominating Set** since  $D = \{1, 3\}$  is a dominating set for  $G$ .

- For a given instance  $(G, k)$  of **Dominating Set** describe a certificate in relation to  $(G, k)$ .
- Provide a semi-formal verifier algorithm that takes as input i) an instance  $(G, k)$ , ii) a certificate for  $(G, k)$  as defined in part a, and decides if the certificate is valid for  $G$ .
- Provide size parameters that may be used to measure the size of an instance of **Dominating Set**.
- Use the size parameters from part c to describe the running time of your verifier from part b. Defend your answer in relation to the algorithm you provided for the verifier.

LO3. Recall the polynomial-time mapping reduction from SAT to 3SAT described in lecture.

- (a) Given the SAT instance  $F(x_1, x_2, x_3) = (\bar{x}_1 \wedge x_2) \vee (x_2 \wedge \bar{x}_3)$ , draw its parse tree and provide the associated Boolean formula  $G$  that is satisfiability equivalent to  $F$  and serves as the beginning step of the reduction. Hint: formula  $G$  introduces  $y$ -variables.
- (b) Rewrite formula  $G$  by making use of the logical identity

$$(P \leftrightarrow Q) \Leftrightarrow [(P \rightarrow Q) \wedge (Q \rightarrow P)]$$

- (c) Rewrite the formula from part b by making use of the logical identity

$$(P \rightarrow Q) \Leftrightarrow (\bar{P} \vee Q).$$

- (d) Rewrite the formula from part c by performing one or more applications of De Morgan's rule.
- (e) Rewrite the formula from part d by performing one or more applications of the distributive rule in order to obtain an AND of OR's. Then convert the AND of OR's to an AND of ternary (i.e. three) OR's and use 3SAT notation to complete the reduction.