NO NOTES, BOOKS, ELECTRONIC DEVICES, OR INTERPERSONAL COMMUNICATION ALLOWED. Submit each solution on a separate sheet of paper.

Problem

LO1. Solve the following problems.

- (a) Show each of the subproblem instances that must be solved when using the recursive multiplication algorithm for finding the product $x \times y$ for x = 29 and y = 57. Make sure to provide the solution to each subproblem instance. Hint: there are seven subproblem instances, including the original problem instance as well as the base case instance with y = 0.
- (b) Consider the RSA key set $(N = 91 = 7 \cdot 13, e = 11)$. Determine the decryption key d.
- LO2. Solve the following problems.
 - (a) Use the Master Theorem to determine the growth of T(n) if it satisfies the recurrence $T(n) = 3T(n/2) + n^{\log_4 16}$. Defend your answer.
 - (b) Use the substitution method to prove that, if T(n) satisfies

$$T(n) = 3T(n/2) + 5n$$

then $T(n) = \mathcal{O}(n^{\log 3})$.

- LO3. Solve each of the following problems.
 - (a) When analyzing a randomized algorithm, what does T(n) represent with respect to the set of random choices made by the algorithm.
 - (b) For the Randomized Quicksort algorithm, provide an interpretation of the recurrence

$$T(n) = T(6) + T(n-7) + O(n).$$

What does it mean and under what assumption(s) is it valid?

(c) Recall that the Minimum Positive Subsequence Sum (MPSS) problem admits a divideand-conquer algorithm that, on input integer array a, requires computing the mpss of any subarray of a that contains both a[n/2-1] and a[n/2] (the end of a_{left} and the beginning of a_{right}). For

$$a = 48, -37, 29, -33, 51, -64, 46, -34, 45, -36$$

provide the two sorted arrays a = LeftSums and b = RightSums from which the minimum positive sum a[i] + b[j] represents the desired mpss (for the middle), where *i* in the index range of *a* and *j* is within the index range of *b*. Also, demonstrate how the minimum positive sum a[i] + b[j] may be computed via the movement of left and right markers.